

1       **Moisture Pathways into the US Intermountain West Associated with Heavy Winter**  
2                               **Precipitation Events**

3  
4       Michael A. Alexander<sup>1</sup>, James D. Scott<sup>1,2</sup>, Dustin Swales<sup>1,2</sup>, Mimi Hughes<sup>1,2</sup>, Kelly Mahoney<sup>1,2</sup>,  
5                               Catherine A. Smith<sup>1,2</sup>

6  
7                               *1 NOAA/Earth System Research Laboratory, Boulder, Colorado*

8                               *2 Cooperative Institute for Research in Environmental Sciences, University of Colorado*  
9                               *Boulder, Boulder, Colorado*

10  
11  
12  
13  
14  
15                               *Submitted to the J. Hydrometeorology*  
16                               *July 2014*  
17  
18  
19  
20  
21  
22  
23  
24  
25  
26  
27  
28  
29

30       Corresponding Author address:

31  
32       Michael Alexander  
33       NOAA/Earth System Research Laboratory  
34       R/PSD1, 325 Broadway  
35       Boulder, CO 80305  
36       E-mail: michael.alexander@noaa.gov

## ABSTRACT

Two methods were used to identify the paths of moisture transport that reach the US intermountain west (IMW) during heavy precipitation events in winter. In the first, the top 150 precipitation events at stations located within six regions in the IMW were identified, then back trajectories were initiated at 6 hour intervals on those days at the four CFSR grid points nearest the stations. The second method identified the leading patterns of integrated water vapor transport (IVT) using the three leading Empirical Orthogonal Functions (EOFs) of IVT over land that were first normalized by the local standard deviation. The top 1% of the associated 6-hourly time series were used to construct composites of IVT, atmospheric circulation and precipitation.

The results from both methods indicate that moisture originating from the Pacific that leads to extreme precipitation in the IMW during winter take distinct pathways through gaps in the Cascade (Oregon-Washington), Sierra-Nevada (California) and Peninsula Mountains (southern California through Baja California). The moisture transported along these routes appears to be the primary source for heavy precipitation for the mountain ranges in the IMW. The synoptic conditions associated with the dominant IVT patterns include a trough ridge couplet at 500 h PA, with the trough located northwest of the ridge where the associated circulation funnels moisture from the west-to-southwest through the mountain gaps and into the IMW.

## 1. Introduction

While generally dry, the interior of the western US can experience extreme precipitation events. For example, in early November 2006, a major rain and flood event occurred in portions of western Montana. One weather station in Glacier National Park recorded over 300 mm (12 inches) of precipitation between 3-8 November and the flow in the North Fork of the Flathead River reached 18,000 cubic feet per second on 8 November exceeding the previous November record by ~40% (Berhardt 2006; Neiman et al. 2008). Extreme precipitation events can also occur over the southwest US during winter, e.g. in late January 2010 a series of three strong storms caused record precipitation and widespread flooding in central Arizona. How does the moisture penetrate the high terrain and complex topography near the US west coast in sufficient amounts to produce extreme precipitation events in the intermountain west (IMW)? Since flow over mountain causes air to cool and thus hold less moisture, air parcels likely take unique pathways through mountain gaps and/or have multiple moisture sources in order to retain enough water vapor to produce intense winter precipitation events in states such as Arizona, Colorado, Idaho and Utah.

Long narrow bands of enhanced water vapor transport generally located ahead of cold fronts, termed atmospheric rivers (ARs, Newell et al. 1992; Zhu and Newell 1998), are associated with heavy rain and snow events along the US west coast (Neiman et al. 2002; Ralph et al. 2004; Ralph et al. 2006, Neiman et al. 2008, 2011; Ralph and Dettinger 2011). Observational studies of strong winter storms over California lead to the identification of key ingredients for heavy precipitation in the Coastal and Sierra Nevada Mountains, including: an ARs transporting warm moist air from the subtropical Pacific (known colloquial as the “Pineapple Express”, McGuirk et al. 1987, 1988; Lackmann and Gyakum 1999), upslope flow

with the atmospheric moisture transport being normal to the topography, a landfalling low-level jet, low static stability, and strong upper-level dynamics (Pandey et al. 1999; Neiman et al. 2002; Ohara et al. 2009). Further inland, the structure and evolution of storms and their associated precipitation are greatly altered by upstream mountain ranges, such as the Cascades and Sierra Nevada. For example, the classical model of a extratropical cyclone with strong near surface fronts often fails in the IMW, where many storms only have elevated baroclinic zones (Marwitz 1980; Schultz et al. 2002).

Recent studies indicate that ARs could move inland and contribute a significant fraction (10%-50%) of wintertime precipitation in the IMW (Leung and Qian 2009; Dettinger et al. 2011; Rutz and Steenburgh 2012). Some pathways for inland ARs have been identified. Rutz and Steenburgh (2012) found that ARs that traverse Baja California contribute significantly to precipitation in Arizona, exemplified by the extreme precipitation/flooding event that occurred in Arizona in January of 2010 (described above) that penetrated through gaps in the mountains along the Baja Peninsula (Neiman et al. 2013 and Hughes et al. 2014). Using combined Empirical Orthogonal Functions (EOFs) of zonal and meridional integrated water vapor transport, Rivera et al. (2014) identified two AR patterns that cause heavy winter precipitation in the Verde River basin in central Arizona, one originates near Hawaii and crosses central Baja California mainly from the west before turning north, and the second originates in the eastern tropical Pacific and crosses southern Baja California from the southwest. Rutz et al. (2014) found that ARs could penetrate into portions of the northwest and into the southwest, including central Arizona, but appeared to be blocked by the high Sierras in California. The Great Salt Lake can also act as a regional source of moisture for heavy precipitation in northern Utah (Steenburgh et al 2000, Steenburgh 2003). While the aforementioned studies have identified some of the moisture



sources for the IMW, the moisture pathways leading to heavy precipitation in the IMW during winter have not been examined systematically.

Air parcel trajectories have been used to identify the sources and transport of moisture, including those associated with ARs. Knippertz and Wernlie (2010) developed a climatology of tropical-to-extratropical trajectories of moist air that identified the Pineapple Express as one of the four main pathways although the trajectories with strong moisture transport that reach the coast of Western North America may also have midlatitude origins (Roberge et al. 2009, Neiman et al. 2013). Using air parcel movements calculated backward in time, or back trajectories, Bao et al. (2006) found that while some of the moisture within ARs that reached the west coast of the United States originated in the tropics much of it resulted from evaporation and moisture convergence in midlatitudes in the warm sector of extratropical cyclones. Back trajectories have also proved useful in understanding precipitation processes and diagnosing moisture sources in other mountainous areas including the Appalachian Mountains (Konrad 1994; Perry et al. 2007; Furman and Konrad 2013).

Extreme precipitation events can cause severe flooding, potentially leading to safety and infrastructure problems. Determining how these events occur in the IMW (taken here to be between the Sierra Nevada/Cascade Mountains and the Continental Divide) would help water managers and emergency planners better prepare for these extreme precipitation events. For example, the Bureau of Reclamation operates hundreds of dams in the IMW and a detailed understanding of the sources of moisture and processes leading to heavy precipitation would inform decisions about dam safety and flood hydrology.

Here we investigate the following questions associated with extreme precipitation events in the US IMW during winter:

- Are there dominant moisture pathways?
- Do they differ for different regions in the IMW?
- Does much of the moisture transport occur through narrow gaps in the mountains, or can moisture remain in the atmosphere after passing over higher topography?
- What synoptic features are associated with the moisture transport?

We will address these questions using two complementary methods: air-parcel trajectories provide a Lagrangian approach for identifying regional pathways into the IMW, while Empirical Orthogonal Function (EOF)/Principal Component (PC) analyses of integrated water vapor transport provide a large-scale Eulerian view of the dominant moisture pathways and the associated synoptic conditions. The precipitation data and atmospheric reanalysis used in this study are described section 2 along with methods used to compute trajectories and EOFs/PCs. The results are presented in section 3 and summarized and discussed in section 4.

## **2. Data and Methods**

We use a combination of data sets and methods to identify the moisture pathways and synoptic conditions associated with heavy precipitation events in the IMW during 1979-2010, the period when high-resolution atmospheric reanalysis is available.

### *a) Station precipitation data*

We obtained daily precipitation data from the Global Historical Climatology Network (<http://www.ncdc.noaa.gov/oa/climate/ghcn-daily/>; Durre et al. 2008, 2010; Menne et al. 2012) to identify heavy precipitation events. We only included stations that had more than 20 years of data where less than 20% of the days were missing observations during 1979-2011 to ensure a

reasonable climatology at each station. The stations were grouped into six regions covering most of the IMW including: 1) eastern Washington, northern Idaho and southeastern British Columbia (hereafter WA-nID) ; 2) eastern Oregon and southern Idaho (OR-sID); 3) Nevada (NV); 4) Utah and western Colorado (UT-CO); 5) southeastern California, western Arizona and southern Nevada (sCA) ; and 6) Arizona and New Mexico (AZ-NM; Fig. 1). While the choice of regions is subjective, their number and boundaries were based on a number of considerations, including: *i*) that regions have more than 50 stations (and all but NV and sCA have more than 100); *ii*) the location of topographic features, e.g. selecting mountain chains for some of the regional boundaries; *iii*) grouping stations that exhibited similar absolute or relative precipitation maxima, e.g. many of the stations in the sCA region exhibited daily precipitation maxima that had a relatively small magnitude but made a substantial contribution to the annual precipitation; and *iv*) the regions with integrated water vapor transport (IVT) maxima based on the EOF analysis (see Section 3.b).

To focus on the heaviest events in our record, we first identified the maximum one-day precipitation amounts during 1979-2011 (~2.5% of winter days) at stations within each of the six regions shown in Fig. 1. The precipitation events are independent in that only the station with the maximum on that day is included. The maximum one-day precipitation values within a region are shown in Fig. 2a via color shading (stations included in the analysis but without a top event are shown by open circles; some stations recorded more than one event). The precipitation amounts over the IMW range from approximately 25-175 mm, with all regions except NV having at least one station with more than 100 mm. The greatest one-day totals (> 125 mm) are found in the central WA-nID, central OR-sID, northern sCA, and western AZ-NM regions.

The percentage of the top 150 events occurring at a particular station in each of the regions is shown in Fig. 2b. Less than 5% of the events occur at most stations in all six regions in the domain. However, the extreme events are not uniformly distributed across a given region: many stations do not have a precipitation event in the top 150, but there are a few stations at which 10% or even 15% of the extremes occur. The concentration of extreme events at just a few stations is most pronounced in the WA-nID, AZ-NM and sCA regions where only three stations account for ~40% of the cases.

#### *b) CFSR*

The primary source for the gridded atmospheric fields used here is the Climate Forecast System Reanalysis (CFSR, Saha et al. 2010) from the National Centers for Environmental Prediction (NCEP). The CFSR is global and includes atmosphere, ocean, land and sea ice models, and an assimilation system to merge the model first guess fields with observations. The resolution of the spectral atmospheric model is T382 (~40 km) with 64 vertical levels, which is nearly an order of magnitude finer than NCEP reanalysis (versions 1 and 2) used in many previous AR studies (e.g. ). The CFSR analysis fields are archived on a 0.5° latitude and longitude grid at 6-hour intervals from January 1979 to March 2011.

#### *c) Livneh precipitation data*

We also use gridded precipitation values from a high-resolution (1/16°) data set recently developed by Livneh et al. (2013) available over the conterminous United States and the Columbia River Watershed in southwest Canada for the years 1915 – 2011. It is an updated version of the Maurer et al. (2002) data set and derived using daily observations from

approximately 20,000 NOAA cooperative observer stations. The daily data are rescaled so that the long-term monthly climatology equals that from the parameter-elevation regressions on independent slopes model (PRISM; Daly et al. 1994). The Livneh dataset likely provides a better representation of precipitation than CFSR in the IMW due to its higher resolution and explicit treatment of topography.

#### *d) Trajectories*

Backward (in time) air parcel trajectories, based on the three-dimensional wind field obtained from CFSR were computed using a method originally developed at the University of Melbourne (<http://www.cycstats.org/trajectories/trajhome.htm>; Noone and Simmonds 1999; Barras and Simmonds 2009), which we modified to provide additional output including the surface pressure. Trajectories were computed for the top 150 one-day precipitation totals (as defined in section 2.a) that occurred at stations within each of the six regions shown in Figs. 1&2. The trajectories were initiated from the four CFSR grid points surrounding the station that recorded extreme precipitation event and were initiated at the four 6-hour intervals on the day the event occurred. In the vertical, we examined trajectories initiated on pressure levels from 600 hPa down to the surface at 50 hPa increments (e.g. 600, 650, 700, etc.). For the results presented here, we used trajectories starting at a single pressure level located between 50 and 100 hPa above (at a lower pressure than) the surface. For example, if the surface pressure was 827 hPa, trajectories were initiated at 750 hPa level. We chose this level, which is generally located in the upper boundary layer/lower free troposphere, because it was high enough so that nearly all (> 99%) of the trajectories remained above the surface over North America but was low enough to be located within the region of strong water vapor transport (e.g., see Neiman et al. 2013, Rivera et al.

2014). A total of 2400 trajectories (150 independent events x 4 CFSR grid points x 4 times per day) were initiated in each region. The position of a trajectory is computed backwards in time over the five previous days at one-hour intervals using the six-hourly three-dimensional CFSR wind fields.

The results from the trajectory analysis will be influenced by several factors, including the location of stations within a region and the frequency with which they get heavy precipitation. The individual trajectories will also be influenced by the resolution of the CFSR, which is relatively fine compared to most reanalyses, but is still coarse (~40 km) relative to the topographic features in the western US. The trajectories are also not tracking moisture directly but represent air parcels initially located near the station recording heavy precipitation; it is possible that some trajectories track a drier sector of a given storm.

#### *e) EOFs of IVT*

We also examine moisture pathways and associated synoptic variability using EOFs of the integrated water vapor transport (IVT). The IVT was derived from CFSR winds ( $\vec{V}$ ) and specific humidity ( $q$ ) on mandatory pressure levels that were vertically integrated from the surface to 300 hPa ( $\sum_{300\text{ hPa}}^{P_{sfc}} \vec{V} q \Delta p / g$ ). The IVT was computed at 6-hr intervals during winter (Oct-Mar) from 1979 to 2011. The mean and standard deviation of IVT over the domain (25°N-60°N, 150°W-105°W) both exhibit a maximum at ~40°N, 145°W over the Pacific that extends northeastward towards the northwest US (Fig. SM1). Given that IVT variability is greater over the Pacific and along the US west coast than in the IMW, we took several steps to highlight IVT patterns in the IMW prior to calculating the EOFs. First, IVT anomalies were computed by subtracting the monthly means from the total field and were then normalized by dividing the anomaly values by

the standard deviation of the 6 hourly data over all years in a given calendar month. Finally, the EOFs were computed using only land points. We examine the three leading EOFs and use their associated PCs to identify when the patterns of IVT and other variables are of large amplitude.

### 3 Results

#### *a. Trajectory analysis*

To illustrate air parcel paths, example backward trajectories were initiated on four different days at the four CFSR grid points surrounding stations that recorded heavy precipitation in the OR-sID region (Fig. 3). While all of the trajectories came from the Pacific, they indicate that moist air parcels can still take diverse routes to reach the lower troposphere over southwest Idaho including central and northern California, Nevada and Oregon (Fig. 3a), central Oregon (Fig. 3b&d), and southern California and Nevada (Fig. 3c). Some of the trajectories indicate slower and/or more circuitous routes, e.g. two loop over northern California, while several others meander over the Snake River Plain in southwest Idaho for trajectories reaching southeast Idaho on 20 Nov 2010 (Fig. 3d). All trajectory segments occur at pressures (color shading) greater than 700 hPa. The trajectories that pass through central Oregon remain at relatively low pressures (approximately 700-800 hPa; relatively high elevations) over the Cascade Mountains, while those that pass through northeastern California are located at higher pressures over California's central valley and remain at relatively high pressure ( $> 800$  hPa) while passing north of the high Sierra Nevada. The trajectories for 11 Mar 1995 illustrate a different path: the trajectories come from the south and stay at relatively low pressure over land; they parallel the high narrow mountain ranges in Nevada and pass through a relatively low point in the mountains in southern California, near  $\sim 35^{\circ}\text{N}$ ,  $117^{\circ}\text{W}$ .

To determine if the air parcels leading to heavy precipitation have preferred routes, the number of hour-long trajectory segments that end within a CFSR grid volume are counted. The counts are performed for the back trajectories for the top 150 events (16 trajectories per event) initiated in each of the six regions and presented as maps of the number of trajectory points throughout a given CFSR column, and as cross sections of the number of counts as a function of height along a transect (indicated by the black line in Fig. 1) spanning the highest portions of the Cascade, Sierra and the Peninsula Mountains (north-south range extending from east of San Diego through Baja California).

Count maps and cross sections for the three northern regions (i.e., WA-nID; OR-sID and NV) are presented in Fig. 4. Broad scale aspects of the topography are also shown on the maps. The count map for the WA-nID region (Fig. 4a) shows the greatest number of trajectory segments in northern Idaho and over southeast Washington, which corresponds with the area where backward trajectories are initiated (Fig. 2). Further west, the axis of higher count values continues along the Washington/Oregon border, over northwest Oregon and the Pacific. While the count map indicates that trajectories can occur across much of the northwest US from central California to the Canadian border (counts > 50), the majority of the trajectories appear to move through the Columbia River Gorge Gap in the Cascade Mountains (as indicated by the break in the 1000 m contours at  $\sim 45.5^{\circ}\text{N}$ ,  $122^{\circ}\text{W}$ ) and continue along the Columbia River valley. Flow through this gorge is supported by the cross section of the number of trajectories originating in WA-nID, in that the greatest number of trajectories pass through the mountains below 800 hPa at  $45^{\circ}\text{N}$  in the v-shaped notch in the topography (Fig. 4b). A modest secondary maxima occurs over central California and passes through an elevated gap in the mountains at  $\sim 41^{\circ}\text{N}$ , to the north of the high Sierras.



The count map and cross section for the OR-sID region, shown in Figs. 4c&d, show a very different pathway than the trajectories that impact northern Idaho. Along the coast the trajectories extend from the Oregon-Washington border to central California but with a preferred path over the Sacramento Valley (northern portion of the central valley), across northeastern California and northwestern Nevada and into southern Washington and Idaho. The map and cross section for this region indicates that the trajectories pass to the north of the high Sierra's through a gap (near Burney, CA) at  $\sim 41^\circ\text{N}$  and over a lower elevation portion of northwest Nevada. Upon reaching Idaho many trajectories follow the Snake River Plain, which has a bowl shape and extends across all of southern Idaho. The counts also suggest a secondary path over the Cascades and across Oregon.

The count map and cross section for the NV region (Fig. 4 e&f) indicate that the highest portion of the Sierras, which extend along a northwest-southeast axis between about  $35^\circ$  and  $38^\circ$  N, are a barrier which most air parcels flow around. One trajectory path is just to the north of the high Sierra, with higher counts over north central Nevada, while a second extends over southern California and into southern Nevada. Downstream of the Sierra Nevada, there are a higher number of counts between the two branches suggesting that the moisture flow may fill in on the leeward side of the mountains. The relatively high elevation (and low pressure) that the air parcels take to reach NV, especially for the northern route, may limit the amount of moisture they carry and contribute to the lack of extreme precipitation in this region.

Maps and cross sections for the three southern regions, UT-CO, sCA and AZ-NM are shown in Fig 5. For all three, the majority of the routes pass over the Pacific and into California or Baja California. The primary axis for the maximum trajectory counts for the UT-CO region (Fig. 5a), is similar to the southern branch for the NV region, i.e. across southern California and Nevada

but now extends into western Utah along the western side of the Wasatch Mountains. A secondary area of high trajectory counts extends along the Utah/Arizona border, where the terrain is relatively low, and continues into southwest Colorado, where there is another maximum near the San Juan Mountains. Some air parcels reaching UT-CO during heavy precipitation events enter the US from the south including the Gulf of California. The accompanying cross-section (Fig. 5b) indicates that the highest number of trajectories pass south of the high Sierras at  $\sim 35^\circ\text{N}$  between 900 and 750 hPa near a gap in the mountains.

The majority of the back trajectories in sCA were initiated and traverse the southern portion of the region (Fig. 5c), which they reach by either utilizing a zonal path near the border with Mexico or by crossing northern Baja California near  $\sim 29^\circ\text{N}$  and then going north over the Gulf of California. The accompanying cross-section (Fig. 5d) indicates that both paths utilize mountain gaps where more trajectories pass through the northern gap.

The trajectory count maximum occurs in the western part of AZ-NM region (Fig. 5e) in the vicinity of the Mogollon Rim, where many extreme precipitation events occurred. The pathways for the AZ-NM are similar to those for the sCA region, except that more trajectories utilize the southern gap at  $\sim 29^\circ\text{N}$  (Fig. 5f), which is wider and deeper than the one near the border.

Count maps of back trajectories over six-hour intervals (e.g. 0-6, 6-12 hours, etc.) are shown for all six regions in the supplementary material. Going forward in time, air parcels extending over a wide swath of the North Pacific propagate to the west/northwest and generally cross the west coast of North America 24-36 hours prior to reaching the target region (Figs. SM2-7).

How unique are these pathways; i.e. are the trajectories on all winter days in general similar to or different than those associated with extreme precipitation events? To address this question we randomly selected days and locations (which could include those with extreme events) to

initialize the back trajectories as follows: for each region we determine the stations at which the top 150 events occurred, then randomly select one of these stations and a date between October and March from 1979-2011 using a random number generator. As in the extreme cases, trajectories were initiated at the four CFSR gridpoints nearest to the station and at four six-hour intervals on the randomly chosen day. This procedure was repeated 300 times based on the number of trajectories needed to obtain stable count results for the random cases; the counts were then scaled by dividing by two so they could be directly compared with the 150 extreme cases. Count maps and cross sections for the randomly generated trajectories in the northern and southern regions are shown in Figs. SM8&9, while the difference in the counts between the extreme and randomly generated trajectories are shown in Figs. 6&7.

The general patterns of the count maps and cross sections for the random generated trajectories (Figs. SM8&9) are similar to those based on the extreme cases in all six regions (Figs. 4&5). For example, as in the extreme cases, a large fraction of the trajectories at randomly selected stations/days for the WA-nID region follow the Columbia River valley passing through the gap in the Cascades. However, consistent differences between the two are apparent as well. The back trajectory counts within the boundaries of all six regions are lower in the extreme compared to the random cases (Figs. 6&7). This suggests faster movement of the air parcels, i.e. fewer trajectories remain in the target region over time, and likely more moisture transport by stronger winds during winter periods with extreme precipitation. In addition, the back trajectories exhibit a more southwesterly path and are generally lower in the atmosphere in the extreme compared to the random cases. For example, in the OR-sID region (Fig. 6c&d), the number of trajectories passing through California and the gap north of the Sierras increases and the number of trajectories through the Columbia Gorge gap decreases in the extreme relative to the random

cases; there are also fewer trajectories on extreme days above (at pressures less than) 700 hPa. One anticipates that air parcels originating further north and/or at higher elevation would be cooler and carry less water vapor. This analysis suggests, that while the pathways into the interior west on days with heavy precipitation might not be unique, strong flow from the southwest at lower elevations is effective at funneling moisture through the mountain gaps.

*b) EOFs and PC based composites*

The three leading EOFs of IVT are shown in Fig. 8a; recall that the EOFs are computed using normalized IVT values over land, so they are better able to depict the patterns of variability in the IMW. The dominant pattern, which explains 28.5% of the variance, has one sign over the western US with a maximum in southern Oregon/Idaho. The 2<sup>nd</sup> EOF, which explains 23.8% of the variance, exhibits a meridional dipole of IVT, with centers over Arizona and along the US-Canada border and the third EOF, which explains 9.9% of the variance, has anomalies of one sign over California and Nevada and anomalies of the opposite sign over the rest of the land points in the domain. The three leading EOFs exhibit broad spatial patterns and explain more than 60% of the variability over the western US.

Histograms of the three leading PCs are shown in Fig. 8b, where the PC indicates the amplitude and phase of the associated EOF pattern over time. The histogram of PC 1, is highly skewed towards positive values, indicating large positive IVT anomalies into the western US, where EOF 1 exhibits maximum variance, are much more common than large negative anomalies. This is consistent with the atmospheric river concept, with a strong narrow band of water vapor transport during storms, and weak/diffuse dry conditions between storms. The second and third PC also exhibit positive skew but not to the extent of PC 1.

EOFs maximize the explained variance over the entire domain and the patterns are required to be orthogonal in both space and time, which can result in nonphysical patterns. This is especially the case for lower order EOFs/PCs. Here the first two leading EOFs, which explain a large fraction of the variance, are separable from the remaining EOFs, while EOF 3 may not be, as the fourth EOF explains 7% of the variance. We examine the synoptic conditions associated with the leading EOFs using PC-based composites of the full anomaly fields but modify the compositing method (described below) to extract realistic synoptic patterns associated with EOF/PC 3.

The EOF analysis was performed using IVT anomalies during all cold season days at 6 hr intervals; since we wish to focus on extreme events, composites were constructed based on when the largest amplitude values ( $> |99\%|$ ) of the three leading PCs occurred. We used the upper 1%, an average of 232 cases, as opposed to the top 150 cases ( $\sim 2.5\%$ ) as done for the precipitation-based trajectory analysis, as here we are using 6 hourly rather than daily values. (The PC-based composites using the top 2.5% of the values <not shown> are similar to the ones presented below albeit with weaker amplitude.) Since EOF 1 covers the entire western US and heavy precipitation is likely associated with large positive IVT PC 1 values, “high” composites are constructed only for periods with large positive PC values. For EOF/PC 2, positive values are associated with strong IVT into the southwest US, while large negative values indicate strong transport over the northern tier states; thus both high and “low” (bottom 1%) composites are constructed for this PC. The pattern identified by EOF 3 suggests the potential for strong IVT anomalies both near the southern and northern border for positive PC 3 values, while negative PC 3 values are associated with positive IVT anomalies in central California. Since the latter would mainly lead

to precipitation in the high Sierras but not in the IMW, we will construct composites using positive PC 3 values.

The synoptic conditions associated with high PC 1 values are illustrated via composite maps (including values over the ocean) of the non-normalized anomalous IVT magnitude, direction, and convergence ( $-\nabla \cdot \text{IVT}$ ), the specific humidity ( $q$ ) at 700 hPa, 500 hPa height, and precipitation from CFSR and the Livneh data set. The composite IVT maps (Fig. 9a&b) show large IVT anomalies extending northwestward from the subtropical Pacific into the northwest US consistent with a strong AR coming onshore in Oregon and California. Smaller-scale IVT anomalies suggest two pathways towards the interior: one through northern Oregon and into Washington, northern Idaho and Montana and a second over central California then extending northeastward across northern Nevada, southern Idaho and extending into Wyoming. The water vapor anomalies (Fig. 9c) emphasize the southern of these two routes and resembles the main trajectory pathway associated with heavy precipitation events in the OR-sID region (*cf.* Fig. 4c). The anomalous circulation exhibits a trough centered west of Vancouver Island and a ridge over the southwest (Fig. 9d). Both the ridge and trough exhibit a southwest to northeast tilt, with a strong gradient between them especially over the northwest US. Assuming geostrophic flow, the height anomalies are consistent with the cyclonic (counterclockwise) IVT vectors west of Canada and anticyclonic (clockwise) IVT west of Baja California with strong onshore flow of moisture into the western US. The anticyclonic IVT center is located west of the 500 hPa ridge due to larger mean values of  $q$  over the Pacific Ocean than the western US, so the winds transport much more water vapor onshore. Lead/lag sea level pressure (SLP) anomaly composites (Fig. SM10) indicate that a low pressure anomaly over the northeast Pacific intensifies and moves northeastward over the two days before reaching a maximum near

Vancouver Island (at zero lag, concurrent with the other composite fields). The SLP anomaly exhibits a more zonal structure and extends further inland than the trough at 500 hPa.

The IVT exhibits strong convergence (Fig. 9b, green shading) over the Sierra and Cascades from California to southern Canada and along the coastal range in northern California. A secondary maximum is clearly evident over mountainous regions of Idaho and western Montana. These results indicate that the moisture transport identified by IVT EOF 1 doesn't just pass through the IMW but converges over the western mountain ranges, with corresponding weak anomalous moisture divergence on the leeward side of some mountains. While a detailed moisture budget analysis is beyond the scope of this study, the convergence of IVT ( $-\nabla \cdot \text{IVT}$ ) is primarily balanced by precipitation minus evaporation plus the time tendency of the column-integrated moisture (e.g., Newman et al 2012 and references therein). Since evaporation and the moisture tendency are generally smaller than the other budget terms during winter over the IMW (e.g. Smith and Kummerow 2013), the IVT convergence and precipitation patterns should be similar. Indeed, the composite maps of precipitation (Fig. 9e&f, note smaller domain relative to other maps) are very similar to those of  $-\nabla \cdot \text{IVT}$  (Fig. 9b). This is consistent with the orographic driver of winter precipitation in the IMW, with moisture transported from the Pacific Ocean into the interior precipitating out when the air impinges on mountain ranges. While the broad pattern of the precipitation anomalies in CFSR and Livneh data sets (Figs. 9e & f, respectively) are similar, fine scale features are much better resolved in the high-resolution Livneh data set and are clearly influenced by the local topography. For example, anomalous heavy precipitation associated with PC 1 occurs near the peaks in the Sawtooth and Salmon River Mountains in central Idaho but the anomalies are negligible just to the south over the Snake River Plain (Fig. 9f).

The PC2 high composite exhibits enhanced moisture transport from the Pacific Ocean, across southern California and the Baja Peninsula, and into the southwestern US (Fig. 10a&b). Within this broad region, there are two enhanced regions of IVT and 700 hPa  $q$ , one along the US/Mexico border and the other across Baja California at approximately 29-30°N (Fig. 10a&c), where the topography is generally less than 800 m, relatively low areas in the Peninsula Mountains. These enhanced areas of IVT transport are consistent with the major paths for trajectories that traverse southern California and Baja California for heavy precipitation events in the three southern regions, especially sCA and AZ-NM (Fig. 4). The circulation anomalies exhibit a wave train with a large trough centered over Oregon/Northern California flanked by ridges to the northwest and southeast, with a strong gradient in the height field from southern California/northern Baja California northeastwards towards Wyoming. The associated SLP anomaly propagates eastward and becomes zonally elongated and centered further east than the 500 mb anomaly (at zero lag) along the border between Idaho, Nevada and Utah (Fig. SM11). The anomalous moisture flux convergence resulting from this flow is extensive (Fig. 10b), with large  $-\nabla \cdot \text{IVT}$  values and heavy precipitation in the Sierra and Peninsula Mountains in California but also in the interior including the Mogollon Rim in Arizona, the Wasatch Mountains in Utah and the San Juan Mountains in southwest Colorado (Fig. 10e&f).

The PC 2 low composite features IVT anomalies with a long fetch across the eastern Pacific extending from ~25°N, 145°W to the Pacific Northwest where it turns and extends eastward along the Canadian border (Fig. 11a&b). A local maximum in IVT and  $q$  extend across Washington, northern Idaho and into Montana (Fig. 11a,b&c), which is consistent with the trajectory path leading to heavy precipitation in the WA-nID region (Fig. 4a). The 500 hPa height pattern features a ridge centered over the IMW and a trough off the Canadian coast, which



are elongated along a southwest-northeast axis (Fig. 9d). As in the previous composites, the anomalous SLP is zonally elongated with a strong gradient along its southern flank, here parallel to the US/Canadian border (Fig. SM12). Strong IVT convergence and heavy precipitation occurs over the Olympic and Cascade Mountains in western Washington and southern British Columbia with a secondary maxima over northern Idaho and southeast British Columbia (Fig 11b,e,f). However, there is anomalous IVT divergence but positive precipitation anomalies in a few narrow meridional bands, such as in central Washington and western Montana extending northwest into Canada. These occur in the lee of mountains and are likely associated with subsidence and drying. This apparent discrepancy with enhanced precipitation in these regions may be due to several factors including: i) enhanced evaporation compensating for the moisture divergence; ii) a non-negligible decrease over time in the column-integrated moisture content; and iii) after assimilation of observations the moisture budget is not balanced in CFSR (or other reanalyses). Weak negative precipitation anomalies also occur over northern California and Colorado.

To determine if EOF 3 represents a coherent pattern we first regressed out the contribution of both EOF 1 and 2 from the full anomaly fields at each time step (the original EOF 3 becomes the leading pattern of the remaining variance). We then computed the correlation between regional averages of the IVT anomalies in the northern center of this EOF (northern Washington; 48°-50°N, 120°-118°W) and the southern center (northwestern Mexico; 28°-30°N, 111°-109°W), which have the same sign (Fig. 8). The correlation between the two regions is -0.07 for all days and -0.25 for the top 1% of the PC values. This suggests that the IVT in the two regions, associated with the original EOF/PC3 are not strongly related but are an artifact of the analysis technique. (In the full data set the two regions are strongly anti-correlated ( $r=-0.83$ ), due to the

out-of-phase relationship between IVT anomalies in these two areas associated with EOF/PC 2). The atmospheric circulation associated with the positive phase of EOF 3 includes a relative small-scale “cut-off” low-pressure center located just west of northern Baja California (not shown). Cutoff lows are fairly common in the wintertime circulation and can cause heavy precipitation in the southwest US (Knippertz and Martin 2007) and are different from the weather patterns associated with the first two EOFs.

The following procedure is used to construct composites that isolate the cutoff lows and the related IVT and circulation anomalies over the southwest. First, based on the full anomaly fields, we identify 6-hr periods with the top 1% of PC 3 values. Next we exclude periods that are in the top or bottom 1% of PC 2 cases, as PC 2 strongly influences IVT in the southwest US. Finally, we retain events where the IVT anomaly is positive in the southern center of EOF 3 located over northwest Mexico. We retain 116 cases that pass these criteria to construct a composite. The composite indicates enhanced IVT transport and  $q$  over southern Baja California, the Gulf of California, northwest Mexico and into southern Arizona and New Mexico (Fig. 12). While the topography in northern Mexico is relatively high, moisture may be able to move through the north-south oriented mountain valleys of northern Mexico and over the low topography along the eastern shore of the Gulf of California. Moisture can also propagate inland from the Gulf of California into Arizona along the Colorado and Gila River valleys. The 500 hPa (Fig. 12d) and SLP field (Fig. SM13) exhibit a cutoff low, just west of northern Baja California with a weak trough to the north, from which the cutoff may have been linked to at a previous time. Over the US, the precipitation exhibits a maximum over the Mogollon Rim in Arizona and western New Mexico. These circulation and precipitation anomalies closely resemble the “Type 2” AR events identified by Rivera et al. (2014). Enhanced precipitation also occurs over western Washington,

likely associate with the weak trough located to the north, perhaps indicating a link between precipitation in the northwest and southwest US identified in the original EOF 3.

#### **4. Summary and Discussion**

Two complementary methods were used to identify the paths that moisture takes to reach the US intermountain west that contribute to heavy precipitation events during winter. In the first, the top 150 precipitation events at stations located within each of six regions in the IMW were identified, then back trajectories were initiated on those days at the four CFSR grid points nearest the stations. While the trajectories track air parcels and not moisture explicitly, by initiating the parcels relatively close to the surface (within 50-100 hPa of the SLP) where the moisture is concentrated, the trajectories should provide a reasonable representation of water vapor movement. The second method identified the leading patterns of integrated water vapor transport, using the three leading EOFs of IVT over land that were first normalized by the local standard deviation. The top 1% of the associated PCs were used to identify 6 hr periods in which the amplitude of the EOF patterns were large and then composites of IVT, atmospheric circulation and precipitation were computed using those periods. To focus on regions where moisture transport was strong, composites were constructed using positive (“high”) values of PCs 1, 2 and 3 and negative (“low”) values of PC2 (with large IVT along the Canadian border); only a subset of these days were used in the PC3-based composites to better isolate realistic synoptic conditions in the southwest US.

The results from both methods clearly indicate that moisture originating from the Pacific that produces extreme precipitation in the IMW US during winter take dominant pathways through gaps in the Cascade (Oregon-Washington), Sierra-Nevada (California) and Peninsula Mountains

(southern California through Baja California). The following paths for different regions were identified (shown schematically in Fig. 13): 1) the Columbia River basin is a conduit for moisture to reach eastern Washington, northern Idaho and western Montana; 2) a surprising path from central and northern California, through a gap to the north of the high Sierras ( $\sim 41^\circ\text{N}$ ), then north into eastern Oregon and Idaho, into the mountains of central Idaho and along the Snake River Plain; 3) through gaps to the north and south ( $\sim 35^\circ\text{N}$ ) of the high Sierras to reach Nevada; 4) through the gap just south of the Sierras into portions of Utah, Colorado and Arizona; and 5) flow through gaps in the Peninsula Mountains near the US-Mexico border, at  $\sim 29^\circ\text{N}$  and over the southern portion of the peninsula that has relatively low topography, bringing moisture to Arizona and western New Mexico. These pathways are consistent with recent studies of the penetration of atmospheric rivers into the IMW; i.e. Rutz and Steenburg (2012) found that the high Sierras strongly block the inland penetration of ARs, while Neiman et al. (2013), Hughes et al. (2014), Rivera et al. (2014), and Rutz et al. (2014), have found that water vapor transport leading to heavy precipitation in the Mogollon Rim passes through low points in the mountains of Baja California. The moisture transported along these routes appears to be the primary source for heavy precipitation for the mountain ranges in the IMW including the Bitterroot Mountains in northern Idaho, Salmon and Sawtooth Mountains in south/central Idaho, the Wasatch Mountains in Utah, the San Juan Mountains in southwest Colorado and the Mogollon Rim in Arizona.

Several factors may influence the moisture pathways identified here. There can be two (or more) pathways to a given region: e.g., air parcels that reach the WA-nID region from California via the mountain gap to the north of the Sierras, in addition to the main pathway along the Columbia River valley. A single pathway can also transverse more than one region; e.g. water vapor transport through the gap south of Sierras can result in heavy precipitation in southeastern

California (sCA), eastern Nevada (NV), the Mogollon Rim (UT-NM), and the Wasatch and San Juan Mountains in southwest Colorado (UT-CO). Also the station locations, which are usually located in valleys likely miss extreme precipitation events at higher elevations and thus the associated pathways. For example, the trajectory analysis did not clearly identify the path that leads to heavy precipitation in the mountains of northern Colorado, such as the Gore or Park Ranges (near  $\sim 40^{\circ}\text{N}$ ,  $106.5^{\circ}\text{W}$ ). Thus, a more quantitative method for identifying regions (e.g., Bernard et al. 2013), including precipitation estimates from higher elevations and using finer resolution data sets that better resolve gaps through the complex topography may lead to a more precise depiction of the moisture pathways in the IMW. In addition, the influence of the seasonal cycle, as well as lower frequency climate phenomena, such as ENSO, on ARs and the moisture pathways into the IMW, warrant further study.

While the topography clearly influences the moisture transport and precipitation in the IMW, it is the large-scale atmospheric circulation and synoptic conditions that ultimately shape the conditions necessary for heavy precipitation to occur. For example, under some conditions the flux of warm moist air may be strong enough so that not all of the moisture precipitates over the Cascade Mountains and can thereby reach eastern Oregon and Idaho (see Figs. 4 and 7). In addition, the pathways identified here are not limited to moist air (Figs. SM8&9). Air also flows through these gaps from the west during dry conditions, albeit more from the north and at higher elevations; e.g. trajectories reaching the OR-sID region often pass through the Columbia river gap.

The general synoptic conditions associated with heavy precipitation events in the IMW include a trough ridge couplet at 500 h Pa, with the trough located northwest of the ridge and a tight gradient between them over and upstream of where the heavy precipitation occurs. The

583 accompanying circulation results in strong moisture transport from the southwest that reach a  
584 maximum near the topographic gaps. While our analyses are based on precipitation and IVT over  
585 the western US, the general circulation features are consistent with studies of ARs that impact  
586 the Pacific coast and feature concentrated moisture transport ahead of a cold front that extends  
587 thousands of kilometers southwestward over the Pacific Ocean. In addition to the large-scale  
588 ridge and trough system in midlatitudes, smaller-scale cutoff lows, which move on shore at  
589  $\sim 30^\circ\text{N}$ , can also transport tropical moisture across southern Baja and the Gulf of California  
590 resulting in heavy precipitation over central Arizona (also see Knippertz and Martin 2007;  
591 Rivera et al. 2014).

592 Determining potential pathways and sources for extreme precipitation events may help  
593 inform decisions about dam safety, flood hydrology, and future monitoring. The trajectory and  
594 EOF/composite analyses performed here could also provide useful indicators for short-term (1-2  
595 day) predictions at dam sites and indicate the attributes of winter storms that generate the  
596 heaviest precipitation in a given region.

## 597 598 **Acknowledgements**

599 We acknowledge the Bureau of Reclamation for their intellectual and financial support for  
600 this project, and in particular thank Levi Brekke, Jason Caldwell, John England and Victoria  
601 Sankovitch. Joseph Barsugli provided helpful suggestions during the course of the project. We  
602 also thank Ian Simmonds from the University of Melbourne for providing the trajectory software  
603 and Ben Livneh at the University of Colorado for providing the gridded precipitation data.

## 604 605 **References**

Bao, J-W., S. A. Michelson, P. J. Neiman, F. M. Ralph, and J. M. Wilczak, 2006: Interpretation of enhanced integrated water vapor bands associated with extratropical cyclones: Their formation and connection to tropical water vapor. *Mon. Wea. Rev.*, **134**, 1063–1080.

Barras, V., and I. Simmonds, 2009: Observation and modeling of stable water isotopes as diagnostics of rainfall dynamics over southeastern Australia *Journal of Geophysical Research*, **114**, D23308, doi:10.1029/2009JD012132, 2009.

Bernhardt, D., 2006: Glacier National Park flooding November 2006. NWS Western Region Tech. Attachment 08-23, 15 pp.

Bernard, E., P. Naveau, M. Vrac, and O. Mestre, 2013: Clustering of Maxima: Spatial Dependencies among Heavy Rainfall in France. *J. Climate*, **26**, 7929–7937.

Daly, C., R. P. Neilson, and D. L. Phillips, 1994: A statistical-topographic model for mapping climatological precipitation over mountainous terrain. *J. Appl. Meteor.*, **33**, 140–158.

Dettinger M. D., F. M. Ralph, T. Das, P. J. Neiman, and D. Cayan, 2011: Atmospheric rivers, floods, and the water resources of California. *Water*, **3**, 445–478. DOI: 10.3390/w3020445.

Durre, I., M. J. Menne, B. E. Gleason, T. G. Houston, and R. S. Vose, 2010: Comprehensive automated quality assurance of daily surface observations. *J. Appl. Meteor. Climatol.*, **49**, 1615-1633.

- Durre, I., M. J. Menne, and R. S. Vose, 2008: Strategies for evaluating quality-control procedures. *J. Appl. Meteor. Climatol*, **47**, 1785-1791.
- Fuhrmann, C. M., C. E. Konrad, 2013: A Trajectory Approach to Analyzing the Ingredients Associated with Heavy Winter Storms in Central North Carolina. *Wea. Forecasting*, **28**, 647–667.doi: <http://dx.doi.org/10.1175/WAF-D-12-00079.1>
- Hughes, M., K. M. Mahoney, P. J. Neiman, B. J. Moore, M. Alexander, and F. M. Ralph: The landfall and inland penetration of a flood-producing atmospheric river in Arizona. Part II: Sensitivity of modeled precipitation to terrain height and atmospheric river orientation. *J. Hydromet.*, in press.
- Knippertz, P., and J. E. Martin, 2007: A Pacific Moisture Conveyor Belt and Its Relationship to a Significant Precipitation Event in the Semiarid Southwestern United States. *Wea. Forecasting*, **22**, 125–144.
- Knippertz, P., and H. Wernli, 2010: A Lagrangian Climatology of Tropical Moisture Exports to the Northern Hemispheric Extratropics. *J. Climate*, **23**, 987–1003.  
doi: <http://dx.doi.org/10.1175/2009JCLI3333.1>



647 Konrad, C. E., 1994: Moisture trajectories associated with heavy rainfall in the Appalachian  
 648 region of the United States. *Physical Geography*, **15**, 227-248. DOI:  
 649 10.1080/02723646.1994.10642514.  
 650  
 651 Lackmann, G. M., J. R. Gyakum, 1999: Heavy Cold-Season Precipitation in the Northwestern  
 652 United States: Synoptic Climatology and an Analysis of the Flood of 17–18 January 1986.  
 653 *Wea. Forecasting*, **14**, 687–700.  
 654  
 655 Lavers, D. A., G. Villarini, R. P. Allan, E. F. Wood, and A. J. Wade, 2012: The detection of  
 656 atmospheric rivers in atmospheric reanalyses and their links to British winter floods and the  
 657 large-scale climatic circulation. *J. Geophys. Res.*, **117**, D20106, doi:10.1029/2012JD018027.  
 658  
 659 Leung, L. R., and Y. Qian, 2009: Atmospheric rivers induced heavy precipitation and flooding in  
 660 the western US simulated by the WRF regional climate model. *Geophys. Res. Lett.*, **36**,  
 661 DOI: 10.1029/2008GL036445.  
 662  
 663 Livneh, B., E. A. Rosenberg, C. Lin, B. Nijssen, V. Mishra, K. M. Andreadis, E. P. Maurer, D. P.  
 664 Lettenmaier, 2013: A Long-Term Hydrologically Based Dataset of Land Surface Fluxes and  
 665 States for the Conterminous United States: Update and Extensions. *J. Climate*, **26**, 9384–  
 666 9392.  
 667  
 668 Marwitz, J. D., 1980: Winter storms over the San Juan Mountains. Part I: Dynamical processes.  
 669 *J. Appl. Meteor.*, **19**, 913–926.

670

671 Maurer, E. P., A. W. Wood, J. C. Adam, D. P. Lettenmaier, and B. Nijssen, 2002: A long-term  
672 hydrologically based dataset of land surface fluxes and states for the conterminous United  
673 States. *J. Climate*, **15**, 3237–3251.

674

McGuirk, J. P., A. H. Thompson, and N. R. Smith, 1987: Moisture bursts over the tropical  
Pacific Ocean. *Mon. Wea. Rev.*, **115**, 787–798.

McGuirk, J. P., A. H. Thompson, and J. R. Schaefer, 1988: An eastern Pacific tropical  
plume. *Mon. Wea. Rev.*, **116**, 2505–2521.

675

676 Menne, M. J., I. Durre, R. S. Vose, B. E. Gleason, T. G. Houston, 2012: An Overview of the  
677 Global Historical Climatology Network-Daily Database. *J. Atmos. Oceanic Technol.*, **29**,  
678 897–910. doi: <http://dx.doi.org/10.1175/JTECH-D-11-00103.1>

679

680 Neiman, P. J., F. M. Ralph, A. B. White, D. E. Kingsmill, P. O. G. Persson, 2002: The Statistical  
681 Relationship between Upslope Flow and Rainfall in California's Coastal Mountains:  
682 Observations during CALJET. *Mon. Wea. Rev.*, **130**, 1468–1492.  
683 doi: [http://dx.doi.org/10.1175/1520-0493\(2002\)130<1468:TSRBUF>2.0.CO;2](http://dx.doi.org/10.1175/1520-0493(2002)130<1468:TSRBUF>2.0.CO;2)

684

685 Neiman, P. J., F. M. Ralph, G. A. Wick, J. D. Lundquist, and M. D. Dettinger, 2008:  
686 Meteorological characteristics and overland precipitation impacts of atmospheric rivers

affecting the West Coast of North America based on eight years of SSM/I satellite observations. *J. Hydrometeorology*, **9**, 22-47.

Neiman, P. J., L. J. Schick, F. M. Ralph, M. Hughes, and G. A. Wick, 2011: Flooding in Western Washington: The Connection to Atmospheric Rivers. *J. Hydrometeorology*, **12**, 1337-1358.

Neiman, P. J., F. M. Ralph, B. J. Moore, M. Hughes; K. M. Mahoney; J. M. Cordeira and M. D. Dettinger, 2013: The Landfall and Inland Penetration of a Flood-Producing Atmospheric River in Arizona. Part 1: Observed Synoptic-Scale, Orographic, and Hydrometeorological Characteristics. *J. Hydrometeorology*, **14**, 460-484.

Newell, R.E., Newell N. E., and C. Scott, 1992: Tropospheric Rivers? A pilot study. *Geophys. Res. Lett.* **19**, 2401-2404. Doi:10.1029/92GL02916.

Newman, M., G. N. Kiladis, K. M. Weickmann, F. M. Ralph, and P. D. Sardeshmukh 2012: Relative contributions of synoptic and low-frequency eddies to time-mean atmospheric moisture transport, including the role of atmospheric rivers. *J. Climate*, **25**, 7341-7361. doi: <http://dx.doi.org/10.1175/JCLI-D-11-00665.1>

Noone, D., and I. Simmonds, 1999: A three-dimensional spherical trajectory algorithm. Research Activities in Atmospheric and Oceanic Modelling. Report No. 28, WMO/TD-No. 942. H. Ritchie, Ed., World Meteorological Organization, 3.26-3.27.

- O'Hara, M. L. Kaplan, S. J. Underwood, 2009: Synoptic Climatological Analyses of Extreme Snowfalls in the Sierra Nevada. *Weather and Forecasting*, **24**, 1610-1624.
- Pandey, G. R., D. R. Cayan, and K. P. Georgakakos, 1999: Precipitation structure in the Sierra Nevada of California during winter. *J. Geophys. Res.*, **104** (D10) 12019–12030.
- Perry, L. B., Charles E. Konrad, T. W. Schmidlin, 2007: Antecedent Upstream Air Trajectories Associated with Northwest Flow Snowfall in the Southern Appalachians. *Wea. Forecasting*, **22**, 334–352.
- Ralph, F. M., and M. D. Dettinger, 2011: Storms, Floods and the Science of Atmospheric Rivers. *EOS, Transactions, Amer. Geophys. Union.*, **92**, 265-266, doi:10.1029/2011EO320001.
- Ralph, F. M., P. J. Neiman, and G. A. Wick, 2004: Satellite and CALJET aircraft observations of atmospheric rivers over the eastern North Pacific Ocean during the winter of 1997/98. *Mon. Wea. Rev.*, **132**, 1721–1745.
- Ralph, F. M., P. J. Neiman, G. A. Wick, S. I. Gutman, M. D. Dettinger, D. R. Cayan, and A. B. White, 2006: Flooding on California's Russian River: Role of atmospheric rivers. *Geophys. Res. Lett.*, **33**, L13801. doi:10.1029/2006GL026689.

729 Rivera, E. R., F. Dominguez, and C. L. Castro, 2014: Atmospheric Rivers and Extreme Cool  
 730 Season Precipitation Events in the Verde River Basin of Arizona. *J. Hydromet.*, **15**, 813-829,  
 731 doi: 10.1175/JHM-D-12-0189.1  
 732  
 733 Roberge, A., J. R. Gyakum, Eyad H. Atallah, 2009: Analysis of Intense Poleward Water Vapor  
 734 Transports into High Latitudes of Western North America. *Wea. Forecasting*, **24**, 1732–  
 735 1747.  
 736  
 737 Rutz, J. J., W. J. Steenburgh, and F. M. Ralph, 2014: Climatological characteristics of  
 738 atmospheric rivers and their inland penetration over the western United States,  
 739 *Mon. Wea. Rev.*, **142**, 905–921. doi: <http://dx.doi.org/10.1175/MWR-D-13-00168.1>.  
 740  
 741 Rutz, J. J., and W. J. Steenburgh, 2012: Quantifying the Role of Atmospheric Rivers in the  
 742 Interior Western United States, *Atmos. Sci. Let.*, **13**, 257-261.  
 743  
 744 Saha, S., and Coauthors, 2010: The NCEP Climate Forecast System Reanalysis. *Bull. Amer.*  
 745 *Meteor. Soc.*, **91**, 1015–1057.  
 746  
 747 Schultz, D. M., and Coauthors, 2002: Understanding Utah Winter Storms: The Intermountain  
 748 Precipitation Experiment. *Bull. Amer. Meteor. Soc.*, **83**, 189–210.  
 749 doi: [http://dx.doi.org/10.1175/1520-0477\(2002\)083<0189:UUWSTI>2.3.CO;2](http://dx.doi.org/10.1175/1520-0477(2002)083<0189:UUWSTI>2.3.CO;2)

750

751 Smith, R. A., and C. D. Kummerow, 2013: A Comparison of in Situ, Reanalysis, and Satellite  
752 Water Budgets over the Upper Colorado River Basin. *J. Hydrometeor*, **14**, 888–905.

753

754 Steenburgh, W. James, 2003: One Hundred Inches in One Hundred Hours: Evolution of a  
755 Wasatch Mountain Winter Storm Cycle. *Wea. Forecasting*, **18**, 1018–1036.

756

757 Steenburgh, W. J., S. F. Halvorson, and D. J. Onton, 2000: Climatology of lake-effect  
758 snowstorms of the Great Salt Lake. *Mon. Wea. Rev.*, **128**, 709–727.

759

760 Zhu, Y. and R. E. Newell, 1998: A proposed algorithm for moisture fluxes from atmospheric  
761 rivers. *Mon. Wea. Rev.*, **126**, 725–735.

762

763

764

## Figure Captions

Fig. 1 a) Topographic map (m, shading, scale on right) of the western United States and Northern Mexico, that shows the six regions (colored text) used to compute the back trajectories: WA-nID (green), OR-sID (dark blue), NV (orange) UT-CO (light blue), sCA(purple), AZ-NM(yellow), mountain ranges are in black, with the major ranges of the Cascades, Sierra's and Rocky Mountains in larger font; River valleys in *italics*; and station locations are indicated by triangles with the color of their respective region. The black curve, extending along the crest of the Cascade, Sierra and Peninsular Mountains, indicates the position of the cross section shown in Figs. (4-7 and SM8&9). B) Cross section of the topography (in hPa) along the black curve shown in a).

Fig. 2. a) The 150 maximum one-day precipitation amounts during 1979-2011 at stations within each of the six regions. Only the station with the maximum on a given day is included, although a station could have more than one event among the 150 cases. Open circles show stations without a top event. b) Percentage of the top 150 events occurring at a particular station in each of the regions.

Fig. 3 Examples of backward trajectories that were initiated near the surface (surface pressure - 50 hPa) at the four CFSR grid points around the station at 00Z on the day during which one of 150 top precipitation events occurred in the OR-sID region. The dates and precipitation amounts are given above the panels for the four cases. The pressure (hPa) along a trajectory segment is shown by the color (blue-red) scale on the lower right and the terrain height (m) by the (green-white) scale on the bottom left.

788

789 Fig. 4. Count Maps (left) and Cross section (right) indicating the number of back trajectories that  
790 pass through a CFSR grid column originating in the a) WA-nID, b) OR-sID, and c) NV regions.  
791 The trajectories are initiated at 00, 6, 12 and 18z on days when one of the top 150 (independent)  
792 precipitation events occurred. The trajectories are initiated at the four CFSR grid point around  
793 the station that recorded the event at a single pressure level located between 50 and 100 hPa  
794 above the surface (see text for more details). A total of 2400 trajectories were initiated in each  
795 region. The position of a trajectory is estimated at one-hour intervals over the five previous days  
796 using the six-hourly three-dimensional CFSR wind fields. Topography is depicted via a 5-point  
797 smoother applied to the 30 arc-second (~1 km) terrain height, with contours at 1000 m (3281 ft),  
798 1500 m (4921 ft) and 2300 m (7546 ft) and stippling above 2300m. Vertical cross section of the  
799 back trajectory counts along the crest of the Cascade, Sierra and Peninsular Mountains (black  
800 curve in Fig. 1) for the a) WA-nID, b) OR-sID, and c) NV regions. The terrain is shown in black.

801

802 Fig. 5. As in Fig 4, but for the count maps (left) and cross sections (right) for the top 150 events  
803 in the UT-CO, sCA and AZ-NM regions.

804

805 Fig. 6. Count maps (left) and cross sections (right) of the difference between the back trajectories  
806 for the top 150 days with precipitation and 300 randomly chosen days (scaled by a factor of 0.5)  
807 over the 1979-2011 winters in the WA-nID, OR-sID, and NV regions.

808

809 Fig. 7. As in Fig. 6., but count maps (left) and cross sections (right) of the difference between the  
810 top and random back trajectories in the UT-CO, sCA and AZ-NM regions.



811

812

813 Fig. 8. a) The three leading EOFs of the 6-hourly CFSR IVT anomalies, normalized by the local  
814 IVT monthly standard deviation, during October-March computed over the western United  
815 States. The anomalies are shown as a regression between the principal component (time series)  
816 and the normalized anomaly. The percent of variance explained by each EOF is shown in the  
817 upper right. b) Histograms of the three leading principal components normalized values.

818

819 Fig. 9. “High” Composites maps constructed from the top 1% (232 6-hourly values) of the first  
820 principal component, (PC 1) of the anomalous: IVT (a) magnitude; (b) IVT direction (vectors)  
821 and IVT convergence/divergence (shading, convergence =  $-\nabla \cdot \text{IVT} > 0$ , mm day<sup>-1</sup>); (c) 700 hPa  
822 specific humidity ( $q$ , g kg<sup>-1</sup>); (d) 500 hPa height (dm); (e) precipitation from CFSR (mm day<sup>-1</sup>);  
823 and (f) precipitation from Livneh (mm day<sup>-1</sup>).

824

825 Fig. 10. As in Fig. 9, but for the “High” Composites maps constructed from the top 1% of PC 2  
826 values.

827

828 Fig. 11. As in Fig. 9, but for the “Low” Composites maps constructed from the bottom  
829 (negative) 1% of PC 2 values.

830


831 Fig. 12. As in Fig. 9, but for the “High” Composites maps based on the top 1% of PC 3 values.

832 To extract a realistic signal from this EOF we also exclude periods that are in the top or bottom  
833 1% of PC 2 cases, and retain events where the IVT anomaly is positive in the southwestern US,  
834 isolating cutoff lows, leaving 116 events in the composite.

835

836 Fig. 13. Schematic (a) map and (b) cross section along the black line in (a) showing the major  
837 moisture pathways from the Pacific Ocean into the intermountain west.

838




**Fig. 1**

839

840 Fig. 1 a) Topographic map (m, shading, scale on right) of the western United States and Northern Mexico,  
 841 that shows the six regions (colored text) used to compute the back trajectories: WA-nID (green), OR-sID  
 842 (dark blue), NV (orange) UT-CO (light blue), sCA(purple), AZ-NM(yellow), mountain ranges are in  
 843 black, with the major ranges of the Cascades, Sierra's and Rocky Mountains in larger font; River valleys  
 844 in *italics*; and station locations are indicated by triangles with the color of their respective region. The  
 845 black curve, extending along the crest of the Cascade, Sierra and Peninsular Mountains, indicates the  
 846 position of the cross section shown in Figs. (4-7 and SM8&9). B) Cross section of the topography (in  
 847 hPa) along the black curve shown in a).

848



849

850

851

852

853 Fig. 2. a) The 150 maximum one-day precipitation amounts during 1979-2011 at stations within each of  
 854 the six regions. Only the station with the maximum on a given day is included, although a station could  
 855 have more than one event among the 150 cases. Open circles show stations without a top event. b)  
 856 Percentage of the top 150 events occurring at a particular station in each of the regions.

857





Fig. 3. Examples of backward trajectories that were initiated near the surface (surface pressure - 50 hPa) at the four CFSR grid points around the station at 00Z on the day during which one of 150 top precipitation events occurred in the OR-sID region. The dates and precipitation amounts are given above the panels for the four cases. The pressure (hPa) along a trajectory segment is shown by the color (blue-red) scale on the lower right and the terrain height (m) by the (green-white) scale on the bottom left.



868

869

870 Fig. 4. Count Maps (left) and Cross section (right) indicating the number of back trajectories that pass  
 871 through a CFSR grid column originating in the a) WA-nID, b) OR-sID, and c) NV regions. The  
 872 trajectories are initiated at 00, 6, 12 and 18z on days when one of the top 150 (independent) precipitation  
 873 events occurred. The trajectories are initiated at the four CFSR grid point around the station that recorded  
 874 the event at a single pressure level located between 50 and 100 hPa above the surface (see text for more  
 875 details). A total of 2400 trajectories were initiated in each region. The position of a trajectory is estimated  
 876 at one-hour intervals over the five previous days using the six-hourly three-dimensional CFSR wind  
 877 fields. Topography is depicted via a 5-point smoother applied to the 30 arc-second (~1 km) terrain  
 878 height, with contours at 1000 m (3281 ft), 1500 m (4921 ft) and 2300 m (7546 ft) and stippling above  
 879 2300m. Vertical cross section of the back trajectory counts along the crest of the Cascade, Sierra and  
 880 Peninsular Mountains (black curve in Fig. 1) for the a) WA-nID, b) OR-sID, and c) NV regions. The  
 881 terrain is shown in black.











Fig. 5. As in Fig 4, but for the count maps (left) and cross sections (right) for the top 150 events in the UT-CO, sCA and AZ-NM regions.




Fig. 6. Count maps (left) and cross sections (right) of the difference between the back trajectories for the top 150 days with precipitation and 300 randomly chosen days (scaled by a factor of 0.5) over the 1979-2011 winters in the WA-nID, OR-sID, and NV regions.







Fig. 7. As in Fig. 6., but count maps (left) and cross sections (right) of the difference between the top and random back trajectories in the UT-CO, sCA and AZ-NM regions.

897



898

899

900

901

902 Fig. 8. a) The three leading EOFs of the 6-hourly CFSR IVT anomalies, normalized by the local IVT  
 903 monthly standard deviation, during October-March computed over the western United States. The  
 904 anomalies are shown as a regression between the principal component (time series) and the normalized  
 905 anomaly. The percent of variance explained by each EOF is shown in the upper right. b) Histograms of  
 906 the three leading principal components normalized values.

907

# CFSR IVT ONDJFM EOF 1 high 1.0% Anom Comp




Fig. 9. “High” Composites maps constructed from the top 1% (232 6-hourly values) of the first principal component, (PC 1) of the anomalous: IVT (a) magnitude; (b) IVT direction (vectors) and IVT convergence/divergence (shading, convergence =  $-\nabla \cdot \text{IVT} > 0$ , mm day<sup>-1</sup>); (c) 700 hPa specific humidity ( $q$ , g kg<sup>-1</sup>); (d) 500 hPa heigh (dm); (e) precipitation from CFSR (mm day<sup>-1</sup>); and (f) precipitation from Livneh (mm day<sup>-1</sup>).

# CFSR IVT ONDJFM EOF 2 high 1.0% Anom Comp




Fig. 10. As in Fig. 9, but for the “High” Composites maps constructed from the top 1% of PC 2 values.

# CFSR IVT ONDJFM EOF 2 Low 1.0% Anom Comp




Fig. 11. As in Fig. 9, but for the “Low” Composites maps constructed from the bottom (negative) 1% of PC 2 values.



# CFSR IVT ONDJFM EOF 3 (modified) high 1.0% Anom Comp





Fig. 12. As in Fig. 9, but for the “High” Composites maps based on the top 1% of PC 3 values. To extract a realistic signal from this EOF we also exclude periods that are in the top or bottom 1% of PC 2 cases, and retain events where the IVT anomaly is positive in the southwestern US, isolating cutoff lows, leaving 116 events in the composite.



**Fig. 13**

Fig. 13. Schematic (a) map and (b) cross section along the black line in (a) showing the major moisture pathways from the Pacific Ocean into the intermountain west.